

AG Carinae: a Luminous Blue Variable with a high rotational velocity¹

J. H. Groh^{1,2}

groh@astro.iag.usp.br

D. J. Hillier²

hillier@pitt.edu

A. Damineli¹

damineli@astro.iag.usp.br

ABSTRACT

We report the detection of broad absorptions due to Si IV $\lambda\lambda 4088\text{--}4116$ in the Luminous Blue Variable (LBV) AG Carinae during its last hot phase (2001-2003). Our NLTE spectral analysis, with the radiative transfer code CMFGEN, revealed the photospheric nature of these lines predicting, however, much narrower and deeper absorption profiles than observed. Using a recently-developed code to compute synthetic spectra in 2D geometry allowing for the effects of rotation, we could match the broad absorptions with a high projected rotational velocity of $190 \pm 30 \text{ km s}^{-1}$ on 2001 April. Analysis of spectra obtained on 2002 March and 2003 January, when the star was cooling, yielded to a projected rotational velocity of $110 \pm 10 \text{ km s}^{-1}$ and $85 \pm 10 \text{ km s}^{-1}$, respectively. The derived rotational velocities are proportional to $R_{\star}^{-1}$, as expected from angular momentum conservation. We discuss the effects of such high rotation on the spectral analysis of AG Car, and on the wind terminal velocity. Our results show direct spectroscopic evidence, for the first time, that a LBV may rotate at a significant fraction of its break-up velocity. Thus, AG Car (and possibly other LBVs) is indeed close to the $\Gamma\Omega$ limit, as predicted by theoretical studies of LBVs.

¹Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, Rua do Matão 1226, Cidade Universitária, 05508-900, São Paulo, SP, Brasil

²Department of Physics and Astronomy, University of Pittsburgh, 3941 O'Hara Street, Pittsburgh, PA, 15260, USA

Subject headings: stars: atmospheres — stars: early-type — stars: individual
(AG Carinae) — stars: rotation

1. Introduction

Massive stars have a tremendous impact on both the chemical and dynamical evolution of a galaxy. They input energy and momentum, and eject their nucleosynthesis products into the interstellar medium (Freyer et al. 2003). A detailed knowledge of massive star evolution is thus crucial to understanding their host galaxies, especially in the early Universe when massive stars are dominant (Matteucci & Calura 2005).

Over the last decade, significant efforts have been made to include the effects of rotation consistently in the evolutionary models for massive stars (Meynet & Maeder 1997). It has been found that the effects of rotation are far from negligible, especially for the most massive stars with high initial rotational velocity. As a consequence of rotation, the evolutionary tracks on the HR diagram, and the duration of the evolutionary stages (main sequence, Luminous Blue Variable – hereafter LBV, and Wolf-Rayet phases), are substantially modified. For a recent review we refer to Maeder & Meynet (2000b).

Rotation is also expected to play an important role in both LBV variability and LBV eruptions (Langer et al. 1999; Maeder & Meynet 2000a; Aerts et al. 2004). Even before reaching the Eddington limit, these stars may lose large amounts of material if they have a significant rotational velocity compared to their critical rotational velocity. Stellar rotation, during the LBV phase, can have an impact on the surrounding circumstellar medium, shaping their ejected nebulae with the characteristic bipolar geometry found around many of these stars (Humphreys & Davidson 1994; Maeder & Desjacques 2001; Dwarkadas & Owocki 2002).

In this Letter we present high-resolution, high S/N optical spectra of the LBV AG Carinae. AG Car has pronounced S Dor type variations, changing its effective temperature from 9,000 to 25,000 K (Leitherer et al. 1994; Stahl et al. 2001). AG Car is considered the prototype of the LBV class and has been extensively observed, from UV to radio wavelengths, with photometric, spectroscopic and polarimetric techniques over several decades. It also has a bipolar nebula which is seen at various wavelengths (Nota et al. 1992; Voors et al.

¹Based on observations made at the 1.52 m ESO telescope at La Silla, Chile and at the 8 m VLT/ESO at Paranal, Chile

2000; Duncan & White 2002) that was ejected approximately 10^4 years ago (Lamers et al. 2001).

Here we report the detection of broad absorptions associated with the high-excitation lines of Si IV $\lambda\lambda 4088\text{--}4116$ in AG Car during its last hot phase (2001-2003). These data are described in Sec. 2, while in Sec. 3 we present the results of our NLTE spectral modeling. Our modeling reveals the photospheric nature of the Si IV lines and allow us to derive, for the first time, the rotational velocity (v_{rot}) of AG Car and how it varies during the S Dor cycle. We show that v_{rot} is high compared to its critical rotational velocity (v_{crit}). In Sec. 4 we discuss some of the effects expected to be present in fast-rotating stars, such as the reduction in the wind terminal velocity and the presence of asymmetric winds. The high v_{rot} derived has important implications not only for AG Car and its wind, but for the general understanding of the LBV phase and the subsequent evolutionary stages of a massive star.

2. Observations

The spectroscopic observations of AG Car analyzed in this work were obtained as part of an observational program to follow the star over a complete S Doradus cycle with a dense time sampling and at high resolution. The results provided by this campaign will be published in a forthcoming paper (Groh et al. in prep.).

During 2001 and 2002, we collected AG Car spectra at the 1.52 m telescope of the European Southern Observatory (ESO) in La Silla, Chile, with the fiber-fed, bench-mounted echelle spectrograph FEROS (Kaufer et al. 1999). To reduce the data we utilized an automatic pipeline reduction system; further details are given in Stahl et al. (1999). The configuration used on the FEROS spectrograph provided a resolution of 48,000 over the region 3600–9200 Å, with increasing signal-to-noise ratio (S/N) from blue to red. We observed AG Car with FEROS on 2001 April 12 and 2002 March 17.

In addition to the FEROS spectra, we retrieved spectra of AG Car from the ESO public-data archive². They were obtained on 2003 January 11 with the UVES spectrograph attached to the 8m VLT-Kueyen at ESO, under ESO Director Discretionary Time Program 266.D-5655 (Bagnulo et al. 2003). The resolution provided by the UVES spectrograph was 80,000 in the wavelength range 3000–10400 Å, with an exceptional S/N of about 400. As with the FEROS spectra, we utilized the pipeline-processed data provided online. Table 1 summarizes the high-resolution spectroscopic observations utilized in this Letter.

²<http://archive.eso.org>

In Figure 1 we show (in velocity space) the absorption profiles of Si IV $\lambda\lambda$ 4088–4116 detected in AG Car during its last hot phase. From bottom to top, we show the 2001 April, 2002 March and 2003 January spectra. The spectra were corrected to the heliocentric frame and the systemic velocity of 10 km s^{-1} (Stahl et al. 2001) was subtracted. The same broad Si IV absorption profiles can also be seen in high-resolution FEROS spectra that we have taken in 2001 January, 2001 June and 2002 July, that are omitted here due to their low S/N, and in lower resolution spectra. Hence, we conclude that the Si IV broad absorptions are present during the whole hot phase (i.e., they are not a transient feature).

3. Modeling and results

To analyze the observational spectra of AG Carinae we used CMFGEN (Hillier & Miller 1998). CMFGEN assumes an outflow in steady-state, spherical symmetry and treats line and continuum formation in NLTE. Each model is specified by the hydrostatic radius R_* , the luminosity L , the mass-loss rate $\dot{M}$, the volume filling-factor f and the chemical abundances Z_i of the included species. The code does not solve the momentum equation of the wind, so a velocity law must be specified. The velocity structure is described by a beta-law modified at depth to match an hydrostatic structure at R_* (typically at a Rosseland depth of 100 in our models), so we also have to input the wind terminal velocity v_∞ and the steepness of the velocity law β . We assumed a constant turbulent velocity of 20 km s^{-1} . A higher turbulent velocity would give a detectable redshift, which is not observed, of the emission lines with high opacity, such the stronger lines of H I and He I.

While CMFGEN does not handle directly the effects of rotation in the radiative transfer, we accounted for its influence on the observed spectrum following the procedures described by Busche & Hillier (2005). For each CMFGEN model we computed the emerging spectrum in 2D geometry viewed from pole (0 degrees, which was equivalent to a non-rotating model), 30 degrees and equator (90 degrees), for a given value of v_{rot} . As our goal is only to show that the broad absorptions are photospheric and broadened by rotation, we did not explore the effects of an equatorial or polar density enhancement on the line profiles. As shown by Busche & Hillier (2005), the first order impact will be on the relative line strengths, which would firstly affect the derivation of the temperature and/or mass-loss rate. These effects will be investigated in a forthcoming paper.

In Figure 2 we show the absorption profiles of Si IV $\lambda\lambda$ 4088–4116 observed on 2001 April compared with our best model. Based on the spectroscopic analysis, we obtained for this epoch $L=1.0 \times 10^6 L_\odot$, $R_*=56 R_\odot$, $T_*=24,340 \text{ K}$, $v_\infty=105 \text{ km s}^{-1}$, $\dot{M}=2.3 \times 10^{-5} M_\odot \text{ yr}^{-1}$, and $f=0.15$. The non-rotating model predicts significantly narrower and deeper absorption

profiles for both Si IV lines than the observations. As these Si IV lines are formed deep in the wind, the broadening effects due to the velocity field are negligible. We needed $v_{rot} \sin i = 190 \pm 30 \text{ km s}^{-1}$ to reproduce the observed lines' shape and width, as can be seen analyzing the fits of model 269_2D_g and 269_2D_c. In spite of the noise of the data, we can clearly rule out a lower value of v_{rot} (e.g. a non-rotating model or model 269_2D_h with $v_{rot} \sin i = 120 \text{ km s}^{-1}$).

To derive v_{crit} , and the surface escape velocity (v_{esc}), we adopted the mass of AG Car as $M = 35 M_{\odot}$ (Smith et al. 1994; Vink & de Koter 2002), which should be regarded as a lower limit (see discussion in Sec. 4 of Vink & de Koter 2002). Changes in M will affect the analysis, since v_{crit} and v_{esc} are both proportional to $M^{0.5}$, and the Eddington parameter Γ depends on M^{-1} . During 2001 April we calculated $v_{crit} = 220 \text{ km s}^{-1}$, following Maeder & Meynet (2000a), for a derived value of $\Gamma = 0.7$. Assuming that we are seeing AG Car edge-on, the derived $v_{rot} = 190 \text{ km s}^{-1}$ amounts to $0.86 v_{crit}$. This value is exceptionally high and it is a lower limit, since higher values will be inferred for lower inclination angles i . For example, if $i = 60^{\circ}$, we would have $v_{rot} = 220 \text{ km s}^{-1}$ and $v_{rot}/v_{crit} = 1$, i.e., the star would rotate nearly at v_{crit} .

The precise determination of v_{rot}/v_{crit} is beyond the scope of this Letter, since it relies on the accuracy of M , L , distance, reddening law, and i . However, for any reasonable change in one or more of these parameters, a significant value of v_{rot}/v_{crit} would still be reached. Hence, the rotationally-broadened photospheric absorptions reported in this Letter make it implausible that AG Car is seen pole-on or from lower inclination angles, as v_{rot} would be arbitrarily high. Moreover, the kinematics and dynamics of its bipolar nebula show extremely low projected velocities, in the range $\pm 50 \text{ km s}^{-1}$ (Nota et al. 1992), which is consistent with a two-lobe polar ejection viewed nearly from the equator.

It is important to point out that our models with rotation predict undetectable changes on the computed profiles of H I, He I lines, and other strong lines on AG Car's spectrum. As they are formed far from the photosphere, the influence of rotation is very low, due to the r^{-1} effect (conservation of angular momentum).

The spectra obtained during 2002 March have a S/N higher than the 2001 April dataset. In Figure 3 we present the best model for this epoch, compared with the observational data. The physical parameters of the star derived from the best fits of CMFGEN were $L = 1.0 \times 10^6 L_{\odot}$, $R_{\star} = 87 R_{\odot}$, $T_{\star} = 19,650 \text{ K}$, $v_{\infty} = 195 \text{ km s}^{-1}$, $\dot{M} = 4.7 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$, and $f = 0.25$. The non-rotating model is inconsistent with the observed lines' width and strength. To match the width and the shape of the lines we had to use a 2D model with $v_{rot} \sin i = 110 \text{ km s}^{-1}$. The high S/N allowed us to constrain the uncertainties to $\pm 10 \text{ km s}^{-1}$. In spite of the superb fits to the line cores and red wings, we can also see in Figure 3 a high-velocity absorption in

the blue wings, which could not be reproduced by our models. Models with higher $\dot{M}$ could not reproduce this feature either.

We present in Figure 4 the 2003 January observations around the Si IV lines compared with our best model. At this epoch AG Car was getting colder, as confirmed by our derived stellar temperature of 17,940 K. Thanks to the high-resolution and high S/N of the VLT data, we could still clearly observe broader Si IV absorptions compared with the predictions of the non-rotating model, with parameters $L=1.0\times10^6 L_\odot$, $R_\star=103 R_\odot$, $T_\star=17,940$ K, $v_\infty=150$ km s $^{-1}$, $\dot{M}=6.0\times10^{-5} M_\odot$ yr $^{-1}$, and $f=0.25$. Running a 2D model with $v_{rot} \sin i=85$ km s $^{-1}$, we could achieve a superb fit to the absorptions' width and strength. A model with $v_{rot} \sin i=100$ km s $^{-1}$ has broader profiles than the observed ones, especially for Si IV $\lambda 4116$. Hence, we could constrain $v_{rot} \sin i=85 \pm 10$ km s $^{-1}$ in this epoch.

In Table 2 we summarize the results obtained for the 3 epochs analyzed in this Letter. The derived luminosity agrees within 20% with previous works (Smith et al. 1994), while the mass-loss rate is clearly variable. A full discussion will be presented in a forthcoming paper. A key result is presented in the last column, where we show the derived value of $v_{rot} R_\star$. This physical parameter should be constant over time due to angular momentum conservation, as the stellar mass is practically invariant on this timescale. Analyzing the values derived in this work, we could verify that the rotational velocities obtained are consistent, taking the errors into account. This reinforces our finding that the broadening detected in the Si IV photospheric lines is a consequence of rotation.

4. Discussion

Rotation is believed to play a significant role during the evolution of massive stars, especially during the short-lived, transitional stages like the LBV phase, where the stellar and wind physical parameters change dramatically on short timescales. Rotation has also been invoked to explain the LBV phenomenon and the origin of asymmetric winds found in some LBVs, like the supermassive star η Car (Smith et al. 2003; van Boekel et al. 2003), and related objects like B[e] stars (Zickgraf et al. 1985). For AG Car, observational evidence for wind asymmetry is provided by polarization data (Leitherer et al. 1994; Schulte-Ladbeck et al. 1994; Davies et al. 2005) and may have a significant impact on the spectroscopic analysis. Such effects are currently being explored by us.

Rotation, and its effects in the wind, have to be taken into account in any theory used to explain how the physical properties change during the S Doradus cycle for AG Car and other LBVs. One of the most noticeable effects of a high rotation, for example, would be a

significant decrease in the wind terminal velocity (Friend & Abbott 1986). Indeed, based on our derived physical parameters, we can calculate that during the hottest epoch (April 2001) $v_{esc}=290\text{ km s}^{-1}$, which includes the reduction due to the radiation pressure. The derived wind terminal velocity of 105 km s^{-1} is incompatible with that surface escape velocity and the predicted value of $v_{\infty}/v_{esc} \sim 1.3 - 2.6$ for hot supergiants (Lamers et al. 1995). However, if we account for the reduction effect that a high rotational velocity has on the effective gravity g_{eff} (Maeder & Meynet 2000a), and hence on the surface escape velocity (defined as $v_{esc} = (2g_{eff} R_{\star})^{0.5}$), we would achieve $v_{esc} \sim 40 - 70\text{ km s}^{-1}$ and $v_{\infty}/v_{esc} \sim 1.5 - 2.6$.

The high rotational velocity also explains why v_{∞} was lower when the star was hotter (2001 April) compared to when the star was cooling (2002 March and 2003 January). At a first glance, this result disagrees with the predicted temperature dependence of v_{∞} for line-driven winds (Vink et al. 2000). However, the effect of rotation on reducing the wind terminal velocity is higher during the hot phase, which can be translated to higher values of v_{rot}/v_{crit} , as shown in Table 2. This can be understood in terms of the $R_{\star}^{-1}$ dependence on the rotational velocity, while the critical rotational velocity depends on $R_{\star}^{-0.5}$.

So far, a high rotational velocity has never been observed directly (i.e. in the stellar spectrum) in any active LBV, in spite of the intense theoretical studies made in recent years. Our spectroscopic analysis shows that AG Car, a well-known prototype of the LBV class, rotates at a significant fraction of its break-up velocity. Interestingly, the behavior of the rotational velocity as a function of the stellar radius during 2001-2003 is consistent with a $R_{\star}^{-1}$ dependence (see Sec. 3), as expected from angular momentum conservation. Our modeling also confirms that AG Car is very luminous ($L=1.0 \times 10^6 L_{\odot}$), and therefore it lies close to the so-called $\Gamma\Omega$ limit (Maeder & Meynet 2000a). Thus, it is likely that rotation, coupled with a high luminosity, plays a key role during the S Dor cycles, the LBV giant eruptions and the subsequent late evolutionary stages of a massive star.

J. H. Groh and A. Damineli thank Brazilian Agencies FAPESP (grant 02/11446-5) and CNPq (grant 200984/2004-7) for financial support. D. J. Hillier gratefully acknowledges partial support for this work from NASA-LTSA grant NAG5-8211. J. H. Groh thanks D. J. Hillier and the University of Pittsburgh for hospitality and support for this work. We thank an anonymous referee for useful suggestions.

REFERENCES

Aerts, C., Lamers, H. J. G. L. M., & Molenberghs, G. 2004, A&A, 418, 639

- Bagnulo, S., Jehin, E., Ledoux, C., et al. 2003, *The Messenger*, 114, 10
- Bjorkman, J. E. & Cassinelli, J. P. 1993, *ApJ*, 409, 429
- Busche, J. R. & Hillier, D. J. 2005, *AJ*, 129, 454
- Davies, B., Oudmaijer, R. D., & Vink, J. S. 2005, *A&A*, 439, 1107
- Duncan, R. A. & White, S. M. 2002, *MNRAS*, 330, 63
- Dwarkadas, V. V. & Owocki, S. P. 2002, *ApJ*, 581, 1337
- Freyer, T., Hensler, G., & Yorke, H. W. 2003, *ApJ*, 594, 888
- Friend, D. B. & Abbott, D. C. 1986, *ApJ*, 311, 701
- Hillier, D. J. & Miller, D. L. 1998, *ApJ*, 496, 407
- Humphreys, R. M. & Davidson, K. 1994, *PASP*, 106, 1025
- Kaufer, A., Stahl, O., Tubbesing, S., et al. 1999, *The Messenger*, 95, 8
- Lamers, H. J. G. L. M. 1997, in *ASP Conf. Ser. 120: Luminous Blue Variables: Massive Stars in Transition*, 76–+
- Lamers, H. J. G. L. M., Nota, A., Panagia, N., Smith, L. J., & Langer, N. 2001, *ApJ*, 551, 764
- Lamers, H. J. G. L. M., Snow, T. P., & Lindholm, D. M. 1995, *ApJ*, 455, 269
- Langer, N., García-Segura, G., & Mac Low, M.-M. 1999, *ApJ*, 520, L49
- Leitherer, C., Allen, R., Altner, B., et al. 1994, *ApJ*, 428, 292
- Maeder, A. & Desjacques, V. 2001, *A&A*, 372, L9
- Maeder, A. & Meynet, G. 2000a, *A&A*, 361, 159
- Maeder, A. & Meynet, G. 2000b, *ARA&A*, 38, 143
- Matteucci, F. & Calura, F. 2005, *MNRAS*, 360, 447
- Meynet, G. & Maeder, A. 1997, *A&A*, 321, 465
- Nota, A., Leitherer, C., Clampin, M., Greenfield, P., & Golimowski, D. A. 1992, *ApJ*, 398, 621

- Owocki, S. P., Cranmer, S. R., & Gayley, K. G. 1996, *ApJ*, 472, L115+
- Schulte-Ladbeck, R. E., Clayton, G. C., Hillier, D. J., Harries, T. J., & Howarth, I. D. 1994, *ApJ*, 429, 846
- Smith, L. J., Crowther, P. A., & Prinja, R. K. 1994, *A&A*, 281, 833
- Smith, N., Davidson, K., Gull, T. R., Ishibashi, K., & Hillier, D. J. 2003, *ApJ*, 586, 432
- Stahl, O., Jankovics, I., Kovács, J., et al. 2001, *A&A*, 375, 54
- Stahl, O., Kaufer, A., & Tubbesing, S. 1999, in *ASP Conf. Ser. 188: Optical and Infrared Spectroscopy of Circumstellar Matter*, p. 331
- van Boekel, R., Kervella, P., Schöller, M., et al. 2003, *A&A*, 410, L37
- Vink, J. S. & de Koter, A. 2002, *A&A*, 393, 543
- Vink, J. S., de Koter, A., & Lamers, H. J. G. L. M. 2000, *A&A*, 362, 295
- Voors, R. H. M., Waters, L. B. F. M., de Koter, A., et al. 2000, *A&A*, 356, 501
- Zickgraf, F.-J., Wolf, B., Stahl, O., Leitherer, C., & Klare, G. 1985, *A&A*, 143, 421

Table 1: Summary of AG Car spectroscopic observations

Epoch	Telescope	Spectral Range ($\text{\AA}$)	R
2001 Apr 12	1.52m ESO	3600-9200	48,000
2002 Mar 17	1.52m ESO	3600-9200	48,000
2003 Jan 11	8m ESO (VLT)	3000-10400	80,000

Table 2: Derived physical parameters of AG Car during 2001-2003, assuming $i = 90^\circ$

Epoch	$T_\star$	$\dot{M}$ ($10^{-5} M_\odot \text{ yr}^{-1}$)	v_∞ (km s^{-1})	v_{rot} (km s^{-1})	v_{rot}/v_{crit}	$R_\star (R_\odot)$	$v_{rot} R_\star$ ($10^4 R_\odot \text{ km s}^{-1}$)
2001 Apr 12	24,340	2.3	105	190 ± 30	0.86	56	10.7 ± 2.2
2002 Mar 17	19,650	4.7	195	110 ± 10	0.63	87	9.5 ± 1.7
2003 Jan 11	17,940	6.0	150	85 ± 10	0.57	103	9.3 ± 1.8

^aWe assume a conservative 10 % error on $R_\star$, accounting for uncertainties on the distance, reddening law and luminosity.




Fig. 1.— High resolution, continuum-normalized spectra of AG Car, centered on Si IV 4088.86 Å (left panel) and on Si IV 4116.10 Å (right panel). The spectra were offset for the sake of clarity. On both panels, we show from top to bottom data from 2003 January (+0.05 offset), 2002 March (no offset) and 2001 April (-0.15 offset). Significant changes in the strength of the Si IV lines, related to changes in the effective temperature of the star, can be seen.




Fig. 2.— High-resolution observed absorption profiles (full line) at Si IV 4088 (left panel) and Si IV 4116 (right panel). A set of CMFGEN models, computed without rotation (dot-dashed), and with a projected rotational velocity of 120 km s^{-1} (dotted line), 180 km s^{-1} (double dot-dashed) and 220 km s^{-1} (dashed line). For the sake of clarity, at the right panel we only show the non-rotating, 180 km s^{-1} and 220 km s^{-1} models. In spite of the errors due to the low S/N, we can clearly rule out a rotational velocity lower than $\sim 150 \text{ km s}^{-1}$ for AG Car on this epoch.




Fig. 3.— Same as Fig. 2, but for the data obtained on 2002 March (full line). The displayed CMFGEN models were computed with (dashed) and without (dot-dashed) a projected rotational velocity of 110 km s^{-1} .




Fig. 4.— Same as Fig. 2, but for the spectra gathered on 2003 January (full line). The CMFGEN models were computed with (dashed) and without (dot-dashed) a projected rotational velocity of 85 km s^{-1} . To estimate the uncertainties, the dotted line shows the CMFGEN model with a projected rotational velocity of 100 km s^{-1} .